



#3

U.S. Department of Commerce Patent and Trademark Office (Form PTO-1449) (Rev. 1-22-91)		Atty. Docket No. 11275/73537	Serial No. 09/031,629
INFORMATION DISCLOSURE STATEMENT BY APPLICANT (Use several sheets if necessary) Page 1 of 1		Applicant: Faustman, D. et al	
		Filing Date: February 27, 1998	Group: 1644

U.S. PATENT DOCUMENTS

Examiner's Initial	Document Number	Date	Name	Class	Subclass	Filing Date if Appropriate

FOREIGN PATENT DOCUMENTS

Examiner's Initial	Document Number	Date	Country	Class	Translation	
					Yes	No
PN	WO 96/13266	stage				
	WO 95/24914	ans				
	WO 95/25533	ans				

OTHER DOCUMENTS (Including Author, Title, Date Pertinent Pages, Etc.)

PN	Aristarkov, et al., 1996, EZ-C, a cyclin-selective ubiquitin carrier protein required for the destruction of mitotic cyclins, <i>Proc. Natl. Acad. Sci. U.S.A.</i> , 93(9): 4294-4299
	Boches, et al., 1982, Role for the Adenosine Triphosphate-Dependent Proteolytic Pathway in Reticulocyte Maturation, <i>Science</i> , 215: 978-980
	Ciechanover, et al., 1984, Ubiquitin Dependence of Selective Protein Degradation Demonstrated in the Mammalian Cell Cycle Mutant ts85, <i>Cell</i> , 37: 57-66
	Driscoll, et al., 1990, The Proteasome (Multicatalytic Protease) Is a Component of the 1500-kDa Proteolytic Complex Which Degrades Ubiquitin-conjugated Proteins, <i>The Journal of Biological Chemistry</i> , 265: 4789-4792
	Eytan, et al., 1989, ATP-dependent incorporation of 20S protease into the 26S complex that degrades proteins conjugated to ubiquitin, <i>Proc. Natl. Acad. Sci. U.S.A.</i> , 86: 7751-7755
	Ganoth, et al., 1988, A Multicomponent System That Degrades Proteins Conjugated to Ubiquitin, <i>The Journal of Biological Chemistry</i> , 263: 12412-12419
	Goldberg, 1992, The mechanism and functions of ATP-dependent proteases in bacterial and animal cells, <i>European Journal of Biochemistry</i> , 203: 9-23
	Goldberg, et al., 1992, Proteolysis, proteasomes and antigen presentation, <i>Nature</i> , 357: 375-379
	Grilli, et al., 1996, Neuroprotection by Aspirin and Sodium Salicylate Through Blockade of NF- κ B Activation, <i>Science</i> , 274: 1383-1385
	Gronostajski, et al., 1985, The ATP Dependence of the Degradation of Short Long-lived Proteins in Growing Fibroblasts, <i>The Journal of Biological Chemistry</i> , 260: 3344-3349

RECEIVED
OCT 09 1998

GROUP 1800



09/031,628

Pn	Haas and Siepmann, 1997, Pathways of ubiquitin conjugation, <i>FASEB Journal</i> , 11: 1257-1268
	Hershko and Ciechanover, 1992, The Ubiquitin System for Protein Degradation, <i>Annual Review of Biochemistry</i> , 61:761-807
	Honda, et al., 1997, Oncoprotein MDM2 is a ubiquitin ligase E3 for tumor suppressor p53, <i>FEBS Letters</i> , 420: 25-27
	Hopkin, Karen, 1997, Regulating Cellular Processes, The Power of Protein Degradation, <i>The Journal of NIH Research</i> , April, 9: 36-42
	Hough, et al., 1987, Purification of Two High Molecular Weight Proteases from Rabbit Reticulocyte Lysate, <i>The Journal of Biological Chemistry</i> , 262: 8303-8313
	Jeffers, et al., 1997, Degradation of the Met Tyrosine Kinase Receptor by the Ubiquitin-Proteasome Pathway, <i>Molecular and Cellular Biology</i> , 17: 799-808
	Kopp and Ghosh, 1994, Inhibition of NF- κ B by Sodium Salicylate and Aspirin, <i>Science</i> , 265: 956-959
	Kwon, et al., 1995, Interleukin-1B-Induced NitricOxide Synthase Expression by Rat Pancreatic B-Cells: Evidence for the Involvement of Nuclear Factor κ B in the Signaling Mechanism, <i>Endocrinology</i> , 136: 4790-4795
	McGuire, et al., 1988, An enzyme related to the high molecular weight multicatalytic proteinase, macroain, participates in a ubiquitin-mediated, ATP-stimulated proteolytic pathway in soluble extracts of BHK 21/C13 fibroblasts, <i>Biochimica et Biophysica Acta</i> , 967: 195-203
	Nocker, et al., 1996, The Multiubiquitin-Chain-Binding Protein Mcb1 Is a Component of the 26S Proteasome in <i>Saccharomyces cerevisiae</i> and Plays a Nonessential, Substrate-Specific Role in Protein Turnover, <i>Molecular and Cellular Biology</i> , 16: 6020-6028
	Orlowski, 1990, The Multicatalytic Proteinase Complex, a Major Extralysosomal Proteolytic System, <i>Biochemistry</i> , 29: 10289-10297
	Rechsteiner et al., 1987, Ubiquitin-Mediated Pathways for Intracellular Proteolysis, <i>Ann. Rev. Cell Biol.</i> , 3: 1-30
	Rivett, 1989, The Multicatalytic Proteinase of Mammalian Cells, <i>Archives of Biochemistry and Biophysics</i> , 268(1): 1-8
	Rivett, 1989, The Multicatalytic Proteinase, <i>The Journal of Biological Chemistry</i> , 264: 12215-12219
	Speiser, et al., 1982, Loss of ATP-dependent Proteolysis with Maturation of Reticulocytes and Erythrocytes, <i>The Journal of Biological Chemistry</i> , 257: 14122-14127
	Tanaka, et al., 1992, Proteasomes: protein and gene structures, <i>New Biol.</i> , 4(3): 173-187
	Townsley, 1997, Dominant-negative cyclin-selective ubiquitin carrier protein E2-C/UbcH10 blocks cells in metaphase, <i>Proc. Natl. Acad. Sci. U.S.A.</i> , 94: 2362-2367
Pn	Waxman, et al., 1987, Demonstration of Two Distinct High Molecular Weight Proteases in Rabbit Reticulocytes, One of Which Degrades Ubiquitin Conjugates, <i>The Journal of Biological Chemistry</i> , 262: 2451-2457
EXAMINER <i>Pattee - J.W.</i>	DATE CONSIDERED <i>10-10-00</i>

EXAMINER: Initial if citation considered, whether or not citation is in conformance with MPEP 609; draw line through citation if not in conformance and not considered. Include copy of this form with next communication